
Chapter 50 Fixed Prosthodontics

Chapter 50 Fixed Prosthodontics

Downloaded from
api.lacavedespapilles.com by Reid &
Lennon

INTRODUCTION TO FIXED PROSTHODONTICS

Fixed prosthodontics is a specialized branch of dentistry dedicated to the restoration and replacement of teeth with permanent, non-removable appliances. Often referred to simply as "fixed dentistry," it encompasses a range of procedures from individual dental crowns to complex full-arch bridges. Unlike removable prostheses (dentures), fixed prostheses are cemented or bonded directly to existing teeth, dental implants, or both, providing stability, durability, and a natural feel. For dental students, the topic "Chapter 50 Fixed Prosthodontics" typically appears in comprehensive textbooks as a deep dive into the planning, design, material science, and clinical execution of these restorations. This article offers an in-depth exploration of fixed prosthodontics, structured to be both informative for practitioners and accessible for students or anyone interested in modern restorative dentistry.

Understanding fixed prosthodontics is essential because these restorations directly impact a patient's ability to chew, speak, and smile with confidence. Advances in digital workflows, adhesive dentistry, and high-strength ceramics have transformed

the field, making fixed prostheses more predictable and esthetic than ever. Whether you are preparing for board examinations, expanding your clinical knowledge, or simply curious about what goes into a dental crown or a fixed bridge, this guide will clarify the principles, materials, and techniques that define Chapter 50 Fixed Prosthodontics.

FUNDAMENTALS OF FIXED PROSTHODONTICS

At its core, fixed prosthodontics involves replacing missing or heavily damaged tooth structure with a prosthesis that is attached to natural teeth or implants. The primary objectives include restoring function, preserving oral health, and achieving esthetic harmony. The discipline requires meticulous treatment planning, an understanding of occlusion, and mastery of dental materials. A fixed prosthesis must withstand significant occlusal forces while maintaining marginal integrity and biocompatibility.

Indications for Fixed

Prosthetic Treatment

Fixed prostheses are indicated in a variety of clinical scenarios. Common reasons include:

- **Extensive tooth decay or fracture** – when a tooth is too damaged for a direct filling but still has a sound root.
- **Endodontically treated teeth** – to protect weakened tooth structure after root canal therapy.
- **Replacing a single missing tooth** – using a fixed bridge or implant-supported crown.
- **Multiple missing teeth** – with conventional fixed bridges, cantilever bridges, or implant-retained prostheses.
- **Esthetic improvement** – such as covering discolored or malformed teeth with ceramic crowns or veneers.
- **Rehabilitation after trauma** – restoring function and appearance following tooth loss or fracture.

Contraindications include poor oral hygiene, active periodontal disease, insufficient tooth structure for retention, or a strong parafunctional habit like bruxism that may lead to restoration fracture.

TYPES OF FIXED PROSTHESES

Dental Crowns

A dental crown, or cap, is a cemented restoration that encases the entire visible portion of a tooth. Crowns are used when a tooth has insufficient clinical crown height or structure to support a filling. They can be made from various materials, including full metal (gold, base metal alloys), porcelain-fused-to-metal (PFM), all-ceramic (feldspathic, lithium disilicate, zirconia), or composite resin. The choice depends on location, esthetic demand, occlusal forces, and cost. For posterior teeth, strength is paramount; for anterior teeth, esthetics often takes precedence.

Fixed Dental Bridges

A fixed bridge replaces one or more missing teeth by using the adjacent teeth (abutments) as supports. A typical three-unit bridge consists of two crowns on the abutment teeth and a pontic (artificial tooth) in the middle. Bridges can be conventional (full coverage), resin-bonded (Maryland bridge), or cantilever. The survival rate of conventional bridges is high, but they require preparation of healthy tooth structure on the abutments, which is a downside. Resin-bonded bridges are more conservative but generally indicated for single missing anterior teeth with minimal occlusal load.

Implant-Supported Prosthesis

Dental implants have revolutionized Chapter 50 Fixed Prosthodontics. Implants are titanium or zirconia posts that osseointegrate with the jawbone, then receive a crown, bridge, or full-arch prosthesis. Implant-supported fixed restorations offer the advantage of maintaining bone volume, avoiding damage to adjacent teeth, and high long-term success rates. Single crowns on implants are now a standard choice for replacing a single missing tooth. For multiple missing teeth, implant-supported bridges or hybrid prostheses provide exceptional stability. The primary challenge is the need for adequate bone quantity and quality, and the higher upfront cost.

MATERIALS IN FIXED PROSTHODONTICS

Material science is central to the success of any fixed restoration. Over the past two decades, the shift toward metal-free restorations has been prominent. Key material categories include:

Metal Alloys

Gold alloys and base metal alloys (cobalt-chrome, nickel-chrome) have exceptional strength and wear resistance, with a long track record of biocompatibility. They are used primarily for posterior

crowns and as substructures for metal-ceramic restorations. Their main disadvantage is poor esthetics, limiting their use in visible areas.

Porcelain-Fused-to-Metal (PFM)

PFM restorations combine a cast metal framework with a layered ceramic veneer. They offer a good balance of strength and appearance, making them a workhorse in fixed prosthodontics for decades. However, the metal core can create opacity and a dark line at the gingival margin. Advances in high-strength ceramics have reduced PFM popularity but they remain a cost-effective and reliable option.

All-Ceramic Materials

All-ceramic systems provide superior translucency and natural color, especially for anterior restorations. Common types include:

- **Lithium disilicate (e.g., e.max)** – high strength, excellent esthetics, suitable for anterior and posterior crowns, bridges up to three units, and implant crowns.
- **Zirconia (tetragonal zirconia polycrystal)** – extremely strong, used for posterior crowns, bridges, and implant abutments. Modern translucent zirconia has improved esthetics, making it popular for anterior use as well.
- **Feldspathic ceramics and leucite-reinforced ceramics**

– used for veneers and inlays/onlays, but less common for full crowns.

Composite Resin

Composite resin crowns are occasionally used for temporary or pediatric restorations, but they lack the wear resistance and color stability of ceramics for long-term fixed prostheses.

CLINICAL PROCEDURES IN FIXED PROSTHODONTICS

The process of delivering a fixed prosthesis involves multiple steps, from diagnosis through final cementation. A typical workflow for a single crown serves as the foundation for understanding other restorations.

Diagnosis and Treatment Planning

This initial phase includes a comprehensive clinical examination, radiographs, diagnostic casts, and sometimes digital impressions using intraoral scanners. The clinician assesses tooth condition, occlusion, periodontal health, and esthetic parameters. A treatment plan is presented, including material choice and cost. For complex cases, a diagnostic wax-up helps visualize the final result.

Tooth Preparation

Tooth preparation is the reduction of the natural tooth structure to create space for the restoration. Principles include:

1. **Preserving sound tooth structure** – avoiding unnecessary removal.
2. **Providing adequate thickness** for the chosen material (e.g., 1.5–2 mm for ceramic, 1–1.5 mm for metal).
3. **Establishing a finish line** (margin) – usually a chamfer or shoulder, placed supragingivally or subgingivally depending on esthetic requirements.
4. **Creating retention and resistance form** – parallelism of axial walls, path of insertion.
5. **Maintaining structural integrity** – avoiding sharp line angles.

For bridges, abutment teeth are prepared in the same manner. Implant-supported crowns require the implant itself to be placed first and then an abutment is connected.

Impression and Provisionalization

Accurate impressions are critical. Traditional methods use elastomeric materials (polyvinyl siloxane) in a tray; digital impressions with intraoral scanners offer convenience and

precision. A provisional (temporary) restoration is fabricated directly or indirectly and cemented temporarily to protect the prepared tooth, maintain occlusion, and preserve gingival contours.

Laboratory Fabrication

The dental laboratory receives the impression or digital file and fabricates the definitive restoration according to the prescription. For ceramic restorations, this may involve CAD/CAM milling, layering of ceramics, and glaze firing. Metal-based restorations require casting or milling. The final restoration is tried in the mouth, checked for fit, contacts, occlusion, and esthetics.

Cementation or Bonding

Cementation is the final step. Depending on the restoration type, different luting agents are used:

- **Conventional cements** (glass ionomer, resin-modified glass ionomer, zinc phosphate) – suitable for metal crowns and well-fitting PFM and zirconia.
- **Adhesive resin cements** – necessary for lithium disilicate, veneers, and areas with compromised retention. These provide micromechanical and chemical bonding to tooth structure and ceramic.

After cementation, the restoration is finished and polished, and occlusion is checked. The patient is given instructions on oral

hygiene and follow-up recall.

COMPLICATIONS AND MAINTENANCE

Fixed prostheses, while durable, can experience complications. Common issues include:

- **Marginal leakage or recurrent caries** – the most frequent cause of failure, often due to inadequate fit or poor oral hygiene.
- **Ceramic fracture or chipping** – especially in older PFM restorations or with bruxism.
- **Loose restoration or debonding** – may occur due to insufficient preparation retention, contamination during cementation, or occlusal overload.
- **Periodontal inflammation** – from overextended margins or plaque accumulation.
- **Loss of tooth vitality** – in some cases, pulpal necrosis can develop after preparation.

Long-term maintenance requires regular dental checkups, professional cleaning, and patient commitment to plaque control around fixed restorations. Floss threaders or interdental brushes are often recommended for cleaning under bridges.

ADVANCES AND FUTURE DIRECTIONS

Chapter 50 Fixed Prosthodontics continues to evolve. Digital dentistry— intraoral scanning, CAD/CAM design, 3D printing of models and temporaries—has improved efficiency and accuracy. Monolithic zirconia with natural translucency is replacing layered

ceramics in many posterior and anterior cases. Implantology is increasingly integrated; guided implant placement and digital prosthetics now allow for same-day teeth. Additionally, biomimetic approaches emphasize preserving dental tissue and using adhesive technologies to mimic natural tooth structure and function.

CONCLUSION

Fixed prosthodontics remains a cornerstone of restorative dental care, offering predictable and long-lasting solutions for damaged

or missing teeth. From the foundational principles taught in textbooks like Chapter 50 Fixed Prosthodontics to the latest digital workflows, the field is rich with options that balance strength, esthetics, and biocompatibility. Whether you are a dental professional refining your skills or a patient considering treatment, understanding the types of prostheses, materials, and procedures can lead to better outcomes. By combining meticulous technique with modern science, fixed prosthodontics restores not only teeth but also the quality of life for countless individuals.